

Some weeds are harder to hate than others. The guavas for instance - they're almost too tasty to condemn. Guava jelly, guava jam, guava shakes.... Read about guava's gastronomic attributes and astronomical spread in two articles in this issue. Ken Langeland and David Hall detail two species (*Psidium guajava* and *P. cattleianum*) in Florida (below) and Charles Wikler describes strawberry guava (*P. cattleianum*) in Hawaii where researchers are hoping to introduce South American biocontrol agents soon.

Find out about the prospects for biocontrol of Chinese privet (*Ligustrum sinense*) in the Southeast on page 17, and saddle up for the wildly-successful "Great Air Potato RoundUp" on page 20. Too bad air potatoes aren't as tasty as guavas. But then again, "Dioscorea shake" doesn't exactly have an appealing ring. -Amy Ferriter

Common guava (*Psidium guajava*)



Exotic Guavas in Florida: so delicious **but—**

By Ken Langeland and
David Hall

The guavas belong to the genus *Psidium* (Myrtaceae, Myrtle Family), which comes from the Greek, meaning edible fruit. *Psidium* is a tropical American genus with approximately 100 species of evergreen trees and shrubs, some with edible berries. Common guava, *Psidium guajava* L., strawberry or Cattle guava, *Psidium*

cattleianum Sabine [*P. littorale* Raddi] (Schroeder 1946, Wunderlin 1998), and Costa Rican guava, *Psidium friedrichsthalianum* (O. Berg) Niedenzu, occur in Florida as landscape plants (Dehgan 1998). *Psidium longipes* (O. Berg.) McVaugh, mangrove berry, is native to Florida. The guava fruit, especially that of *Psidium guajava* cultivars that have been developed with superior taste qualities, is highly regarded for its edible qualities, eaten

fresh and used in jellies, jams, drinks, pies, ice cream, etc. Cultivars of *P. guajava* are grown commercially for their fruit. Common guava, and strawberry guava are known to have escaped cultivation and included on the "Florida Exotic Pest Plant Council's (FLEPPC) 1999 List of Invasive Species". Both are listed under Category I, which means they are considered by FLEPPC as invasive exotics that are altering native plant communities.



Strawberry guava has characteristic gray-brown to reddish bark that peels off in irregular patterns.

Common Guava

Common guava (*Psidium guajava* L.) is a shrub or small tree, usually under 15 feet tall (rarely to 30 feet), with scaly bark. Branches are 4-angled and hairy. The opposite leaves have a short stalk/petiole and elliptic to oblong leathery blades. The lower blade surface is covered with soft hairs, contains numerous glandular dots and has very noticeable riblike lateral veins. Inflorescences are usually 1-flowered in leaf axils. The five sepals are united. The five petals are white, showy, rounded and just under 1 inch long. Stamens are numerous. The ovary is imbedded below the rest of the flower parts (inferior). The usually dull yellow fruits are rounded or pear-shaped, manyseeded berries about 1 to 2 1/2 inches long with pink or yellowish flesh. Reproduction is by seeds. Blooming occurs all year.

Common guava has been cultivated and distributed by man and animals for so long that it is difficult to be certain of its place of origin but it is believed to be from an area extending from southern Mexico into or through Central America (Morton 1987). It is now cultivated and escaped throughout the New and Old World tropics. In addition to its value for edible fruit, the wood is valued in some parts of the world, tannin is extracted for use in tanning and for dyes and, in addi

Psidium cattleianum, Deliciously dangerous in Hawaii

By Charles Wikler

Eating the delicious red or yellow fruits of strawberry guava (*Psidium cattleianum*) we would never imagine how big the problem caused by these plants is in many tropical regions. This species is invading extensive areas, dispersing into important natural habitats, ecologically disrupting them. In addition, this plant is host to several species of fruit flies.

Introduced to Hawaii about 1825, strawberry guava quickly escaped from cultivation. It has become the most important forest weed in Hawaii (Smith, 1985) due to suitable soil and climatic factors, the absence of natural enemies, and effective dispersal by several different agents. Initially, birds disseminated seeds. Feral pigs, relatively recent invaders of the Hawaiian ecosystem, have since become the most important dispersal agent of strawberry guava seeds (Diong, 1982).

Strawberry guava possesses several of the characteristics of an opportunist invasive species, but it seems that its ability to grow clonally is the main reason for its successful invasion and domination of intact native forests in the Hawaiian archipelago (Huenneke, 1991). This plant is an excellent example of a plant

once confined to a small region but now a widespread problem throughout the tropics and subtropics.

From the beginning, studies about this plant generated considerable debate over which scientific name was published correctly for *Psidium cattleianum*. Sabine (1821) named the plant in honor of William Cattley, the first person to successfully cultivate the species in Britain. Some authors considered Raddi as the first to describe the plant, also in 1821, but in the literature search, Raddi's description refers to the plant as *Psidium littorale* in 1823. Thus, the name *Psidium littorale* is usually considered a junior synonym, as the original descriptions of both represent the same species.

According to Fosberg (1941) in Ellshoff et al. (1995), the common name used for strawberry guava is "araca" in Brazil, "waiwai" in Hawaii, and "Cattley guava" among some American horticulturists. Strawberry guava is also mentioned in the literature as "goiave de L'Afrique", in the Dominican Republic, "araca-saiyu" and "guayabo amarillo" in Argentina, "araza" in Uruguay, "Calcutta-guava" in India, "china-guava" for the British,

"goyavier of St. Martin" in Guadeloupe, "goyavier fraise" for the French, "goyavier prune" in Martinique and "purple-guava" in Jamaica.

Hoehne (1946) mentioned that the diKerentiahon between the species becomes complicated for what is called "araca." Although the exact origin of the type specimen is not known, it seems that the plant was cultivated in Asia and America. However, the type locality of *P. littorale* is cited by Raddi (1823) as the coasts of Brazil.

Two botanical forms of strawberry guava tree are recognized based on fruit color. They are the red fruit *P. cattleianum* f. *cattleianum* (*P. littorale* var. *longipes*) and the yellow fruit *P. cattleianum* f. *lucidum* (*P. littorale* var. *lucidum*).

Geographical distribution in the area of origin

In Brazil, the genus *Psidium* is represented by 9 species, including *P. cattleianum*. The plant occurs from the South of Espirito Santo State in Brazil to Uruguay, (from 20° and 32° S). Strawberry guava is a characteristic bush of the "restinga," typical vegetation of the Atlantic Rain Forest, where it is widely

dispersed. Although present, it is not too frequent in the first plateau. The species occurs in humid soils, in small woods (capoeiras), and the border of creeks and semi-devastated forests. It is also quite frequent in the coastal swampy fields along the littoral zone of the southern states of Santa Catarina and Rio Grande do Sul. It can be found in and around the southern plateau, occurring in the ciliary forests, as well as in disturbed fields (Reitz, Klein & Reis, 1983).

Morpholog of the varieties of *Psidium cattleianum*

According to Wikler (1999), the main differences between the yellow and red forms of strawberry guava are:

- *P. cattleianum* var. *lucidum* (yellow form)

It is commonly a shrub in the coast, as part of a vegetation formation called "restinga", where a wide variety of plants grow on the sandy substrate with a high water table. The climate in the coast is hot and humid with temperatures between 18 and 22°C and annual rainfall between 1430-2450 mm. Its height varies between 2.5 and 4 meters, rarely exceeding 5 m high. The trunk is tortuous and with very characteristic grayish-brown coloration, being

unmistakable in the places where it is found. Its crown is round and quite dense. The fruits are yellow, as well as the endocarp that ranges from light yellow to white. In Parana State it is distributed along the coast, and on the first and second plateaus.



Psidium cattleianum var. *lucidum*

- *P. cattleianum* var. *cattleianum* (red form)

The red form is a tree, with heights from 2.5 meters up to 20 meters. Its trunk has the same color of the yellow form although it differs in shape being straight and cylindrical with a larger diameter. Its crown is slightly elongated and quite dense. Its fruit and its endocarp are red colored. In the Araucaria Forest it is found inside the woods but exclusively in the first plateau, ranging from 6501100 m. The climate is hot and humid, with temperature between 15 - 19 °C and 1250 - 2500 mm annual rainfall.



Psidium cattleianum var. *cattleianum*

Biocontrol in Hawaii

Unable to control strawberry guava by mechanical and chemical means in Hawaii, the U.S. National Park Service initiated efforts to find biological control agents against this weed. A cooperative program between the Cooperative National Park Resources Studies Unit, University of Hawaii at Manoa, and the Parana Forest Research Foundation and the Federal University of Parana was established in March 1991.

S O M E

EXOTIC

SPECIES NEED TO

RUN WILD

Others Don't. Improved JLB Oil Plus makes your basal-applied herbicides work better. Better penetration. Better uptake. A natural vegetable oil based diluent at a reasonable cost. Order today from "The Adjuvant Experts."

HELPS CONTROL:

- Brazilian Pepper
- Chinese Tallow Tree
- Australian Pine
- Downy Rose Myrtle
- Salt Cedar (Tamarisk)

800-228-1833

BREWER International

P.O. BOX 6006 - VERO BEACH, FL 32961-6006 - WWW.BREWERINT.COM - FAX: 561-778-2490

tion to other medicinal uses, the roots, bark, leaves and immature fruits are used to halt gastroenteritis, diarrhea and dysentery throughout the tropics (Morton 1987). In Mexico, the tree may be parasitized by mistletoe that causes rosette-like formations called wood flowers, which are sold as ornamental curiosities (Morton 1987).

According to P. W. Reasoner, common guava was introduced to Florida from Cuba in 1847 (Popenoe 1920). However, it was already reported as naturalized in 1765 (DeBrahm 1773). The 1887-1888 Catalog and Price List for Royal Palm Nurseries states: "The guava has become a necessity to South Florida; is to South Florida what the peach is to Georgia". It has been grown successfully as far north as the Pinellas peninsula [sic] on the west coast and Cape Canaveral on the east coast (Popenoe 1920) and was grown commercially at Palma Sola, Punta Gorda, Opalacha, Indiantown, and other localities (Barrett 1956). Three cultivars are now grown commercially, Homestead and two private selections. About 225 acres are in production in Dade (90%), Lee, Broward, and Palm Beach Counties (Jonathan Crane, personal communication). This acreage has been expanding since 1992 and has greater potential due to expanding Asian and Latin markets. The crop is worth in excess of \$3,000,000 annually. In Hawaii, 7,000 acres are grown. While available in the horticultural trade, common guava is relatively insignificant in the current ornamental grower and landscape market and a phase-out of production and sale by Florida ornamental growers is encouraged by an agreement between the FLEPPC and the Florida Nurserymen and Growers Association (Aylsworth 1999).

Common guava grows and fruits under unfavorable conditions, spreads rapidly by seeds and has become a weed in many areas (Popenoe 1920). In fact, it is present as a weed in 27 countries and considered a common to serious pest in nine of these (Holm et al. 1979). It forms thickets and has a serious impact on native forests and open woodlands (Cronk and Fuller 1995). Small, in 1933, reported it in hammocks, roadsides, pinelands, and

old fields of peninsular Florida and the Keys, and it was further reported spreading into hammocks and pinelands in 1971 (Long and Lakela). It now ranges from Pinellas and Brevard Counties south to the Keys (Nelson 1994) and has been reported from Florida parks and preserves in Broward, Collier, Dade, Highlands, Lee, Martin, Osceola, Palm Beach, and Sarasota Counties (FLEPPC 2000). In Jonathon Dickinson and Seabranck State Parks alone 1,163 stems were treated in 1998-99 (Florida Park Service). Trees are controlled by basal bark applications of Garlon 4, cut stump applications of Garlon 4 or 3A, and hand pulling of seedlings.

Strawberry Guava

Strawberry guava [*P. cattleianum* Sabine] is a shrub or small tree to 25 feet tall with smooth, gray-brown to reddish bark that is inclined to peel off in irregular patterns. The slick, leathery, evergreen, opposite leaves are elliptic to somewhat broader towards the tip and up to 3 inches long with a short stalk/petiole. The lateral nerves in the blades are ascending and curve forward without forming a distinct marginal nerve. The inflorescence is stalked and 1-flowered from the leaf axils. Flowers have five united sepals. The five petals are white, showy, rounded and about 1 inch wide. Stamens are numerous and anthers are attached in the center and split longitudinally. The ovary is imbedded below the rest of the flower parts (inferior). The red-purple or yellow fruits are almost

A survey identified potential biological control agents from which the impacts of seven insect species and their effects on *Psidium cattleianum* are discussed by Wickler et al. (1999): a lead gall produced by *Tectococcus ovatus* (Homoptera, Eriococcidae), bud galls formed in response to *Dasineura gigantea* (Diptera, Cecidomyiidae), a shoot gall produced by *Eurytoma* sp. (Hymenoptera, Eurytomidae), a seed gall induced by *Sycophilia* sp. (Hymenoptera, Eurytomidae), another leaf gall formed in response to an unknown species of Psyllidae, the sawfly, *Haplostegus epimelas* (Hymenoptera, Pergidae), and the chrysomelid *Lamprosoma azureum* (Coleoptera, Chrysomelidae).

The author gratefully acknowledges Dr. Clifford W. Smith for all his assistance since the beginning of these studies, Dr. Simon Elliot and Milton Mendonça Jr. for their helpful comments in the draft manuscript and Dr. J.H. Pedrosa-Macedo and the Biological Control of Weeds group from the Laboratory of Forest Protection of UFPR for their further assistance. Funding from the U.S. Na-

Solution

WATER SOLUBLE® AMINE

FROM

Riverdale

**User & Environmentally Friendly
Water Soluble 2,4-D Amine
In Convenient Water Soluble Bags.**

**For The Industrial Vegetation Manager
Concerned About Applicator Safety & Exposure**

- No Measuring • No Pouring • No Freezing
- No Exposure To Concentrates • Low Odor
 - Totally Soluble • Won't Settle Out
- No Triple Rinsing • No Container Disposal
 - No WPS Label Language
- Packaged In Recyclable Cartons
 - Contains Full Aquatic Label

**for more information, call
(800)345-3330**

round, many-seeded berries about 1 and 1/2 inches long with white flesh. Reproduction is by seeds. Blooming usually occurs in early summer. Native to southeastern Brazil, it is planted in the subtropics and tropics. Strawberry guava can survive temperatures as low as 22 F (Morton 1987) and is grown as far north as Alachua County.

P. cattleianum occurs as two distinct botanical varieties. The common redfruited variety is called Cattley guava or strawberry guava, *P. cattleianum* Sabine var. *cattleianum* [*P. littorale* Raddi var. *longipes* (O. Berg) Fosberg] (Fosberg 1941). This variety is a small dense tree. It makes a nice ornamental with the copious array of dark shiny green leaves and bright red fruits. The fruit is sweet and obviously eaten and spread by wildlife and cattle. As an escape, strawberry guava has been found on floodplains and in oak hammocks and cattle pens from Osceola County south into Lee County (Wunderlin *et al.* 1995). The common name, strawberry guava, comes from the flavor of the fruit, which tastes like a strawberry. The fruits are eaten fresh or used for making jelly, drinks and other culinary uses, however, it is not grown commercially for its fruit in Florida. It is an attractive dooryard tree, presenting "_____ an almost exotic or Japanese picture as an accent" and used as a hedge (Stressau 1986).

The less frequent yellow-fruited variety is called yellow Cattley guava or yellow strawberry guava, *P. cattleianum* Sabine var. *lucidum* Hort. [*P. littorale* Raddi var. *lucidum* (Degener) Fosberg] (Fosberg 1941). This variety is a loosely branched small tree said to be useful for reforestation (Staff, Hortus Third 1976). The fruit is sulphur-yellow, translucent and somewhat acid when ripe. Yellow Strawberry guava forms dense thickets along the margin of the Little Manatee River in Hillsborough County. This variety has also been listed [*sic*] as *P. lucidum* Hort., *P. chinense* Lodd. ex Lodd. and *P. sinense* (not a valid name, probably a misspelling of *chinense*) (Popenoe 1920).

Strawberry guava forms thickets and shades out native vegetation in forests and open woodlands (Cronk and Fuller 1995). It has had cata-

strophic effects on native habitats of Mauritius, and it is considered among the worst pest plants in Hawaii (see Wikler, this issue), where it has invaded a variety of natural areas (Cronk and Fuller 1995). It has become dominant in native forests of Hawaii, including in two national parks, where its clonal spread is enhanced by activities of feral pigs (Huenneke and Vitousek 1990).

Strawberry guava first appeared in the horticulture trade in Florida in the 1887-1888 Catalog and Price List for Royal Palm Nurseries. However, it was not included in southeastern flora by Small (1933). It was noted as "often growing wild" in 1956 by Barrett (1956). It has been reported in Florida parks and preserves in Pinellas, Hillsborough, Martin, and Palm Beach Counties (FLEPPC 1996). In Jonathan Dickinson State Park and Seabranck Preserve alone, 2,087 Strawberry guava stems were controlled in 1998-99 (Florida Park Service). Where possible, stems are cut and treated with Garlon 3A. In Jonathan Dickinson State Park, it was necessary for DEP to bulldoze and restore a three acre site (Richard Roberts, personal communication). At Seabranck, up to seven guava plants per square meter occur and 5-15% of vegetation has consisted of Strawberry guava in various sites in 35 acres John Griner, personal communication). Twenty acres have been restored by selective removal of this pest plant and planting of native vegetation.

In summary, the guavas have provided us with delicious desserts, ornamental plantings, and commercial income. On the other hand, common and Strawberry guavas have escaped cultivation and are weeds in natural areas requiring expenditure of public funds to protect native plant communities. This is an example of the dilemma we face with many introduced plants that are both commercially valuable and natural area weeds.

Acknowledgements

Thanks to Hillary Cherry for library assistance and Bob Pemberton for supplying information from the Royal Palm Nursery catalogue.

Ken Langeland is a Professor at the University of Florida's Center for Aquatic

tional Park Services and the University of Hawaii (CA80xx2-9004) supported this study.

Charles Wikler is a Forester at the Paraná Forest Research Foundation and holds an M.S. and Ph.D. in Biological Control of Insects and Weeds Rua Theresa P. Moura, 70 - Pilarzinho 82100-440 - Curitiba - Paraná - Brazil. E-mail: cwikler@bbs2.sul.com.br

References

- Diong, C.H. 1982. Population biology and management of the feral pig (*Sus scrofa*) in Kipahulu Valley, Maui. Unpublished Ph.D. dissertation, University of Hawaii, Honolulu.
- Ellshoff, Z.E. Gardner, D.E., Wikler, C., & Smith C.W. 1995. Annotated bibliography of the genus *Psidium*, with emphasis of *P. cattleianum* (strawberry guava) and *P. guajava* (common guava), forest weeds in Hawai'i. *Technical Report No. 95*. Cooperative National Park Resources Studies Unit, Botany Dept., University of Hawaii, Honolulu.
- Hoehne, F.C.; 1946. Frutas Indígenas. Instituto de Botânica, Publicação da série "D", São Paulo. 88 p.
- Huenneke, L.F. 1991. Population biology of an invading tree, *Psidium cattleianum*, in Hawai'i Volcanoes National Park. Pp. 177-188, In: Center, T.D., R.F. Doren, R.L.
- Hofstetter, R.L. Meyers, and L.D. Whiteaker (eds.) Proceedings of the Symposium on Exotic Pest Plants; U.S. Department of the Interior, National Park Service; November 2-4, 1988; Miami, Florida.
- Reitz, P.R., Klein, R.M. & Reis, A. 1983. Flora Catarinense (*Psidium*). (Flora of Santa Catarina (*Psidium*)). *Sellowia* 35:684-715.
- Sabine, J. 1821. *Psidium cattleianum*. *Trans. Roy. Hort. Soc.* 4:315. Pl. 11.
- Smith, C.W. 1985. Impact of alien plants on Hawaii's native biota. In: Stone, C.O., and J.M. Scott (eds.), *Hawaii's Terrestrial Ecosystems: Preservation and Management*. Cooperative National Park Resources Studies Unit, University of Hawaii, Honolulu.
- Wikler, C. 1999. Distribuição geográfica mundial de *Psidium cattleianum* Sabine (Myrtaceae) e um cecidógeno com possibilidades de utilização em controle biológico.
- Post-Graduation Forestry Course. Universidade Federal do Paraná. Doctoral thesis. 135 p.
- Wikler, C., Pedrosa-Macedo, J.H., Vitorino, M.D., Caxambú, M.G. & Smith, C.W. 1999. Strawberry Guava (*Psidium cattleianum*) – Prospects for Biological Control. Program and abstracts of the X International Symposium on Biological Control of Weeds, P128, pp. 138. N. Spencer and R. Nowierski (eds.) 04-14 July, Bozeman, Montana, USA. Montana State University.

and *Invasive Plants in Gainesville, FL*.
EMail: kal@gnv.ifas.ufl.edu

David W. Hall is President of Hall Consultant, Inc. in Gainesville, FL.

References

- Aylsworth, J.D. 1999. Invasive is Out. Ornamental Outlook. 8(8):40,42.
- Barrett, M.F. 1956. Common Exotic Trees of South Florida. Univ. of Florida Press, Gainesville. pp. 298-299.
- Florida Park Service. Undated. 1998-99 Resource Management Annual Report - July 1, 1998-June 30, 1999.
- Cronk, Q.C.B., and J.L. Fuller. 1995. Plant Invasions. Chapman and Hall, London. pp. 111-114.
- DeBrahm, W.G. 1773. DeBrahm's Report of the General Survey in the Southern District of North America. Tricentennial edition (L. DeVorse, Jr., ed.), No. 3, 1971. Univ. of So. Carolina Press, Columbia. 325 pp. As cited in: Austin, D.E. 1978. Exotic plants and their effects in southeastern Florida. Env. Conserv. 5:25-34.
- Dehgan, B. 1998. Landscape Plants for Subtropical Climates. Univ. Press of Florida, Gainesville. p. 480.
- FLEPPC. 2000. Florida Exotic Pest Plant Council occurrence data base. Unpublished. Data available by requests via Web site: <http://www.fleppc.org/>.
- Fosberg, ER. 1941. Varieties of the Strawberry

- Guava. Proc. Biol. Soc. Wash. 54:179-180.
- Holm, L., J.V. Pancho, J.R Herberger, and D.L. Plucknett. 1979. A Geographical Atlas of World Weeds. John Wiley & Sons, New York. pp. 297-298.
- Huenneke, L.E, and P.M. Vitousek. 1990. Seedling and clonal recruitment of the invasive tree *Psidium cattleianum*: Implications for management of native Hawaiian forests. Biol. Cons. 53:199-211.
- Long, R.W., and O. Lakela. 1971. A Flora of Tropical Florida. University of Miami Press. Coral Gables, FL. pp. 642-643
- Morton J.E 1987. Fruits of Warm Climates. Published by the author, Miami, FL. pp.356-367.
- Nelson, G. 1994. The Trees of Florida. Pineapple Press Inc., Sarasota, FL. pp. 104-105, 241-242.
- Popenoe, W. 1920. Manual of Tropical and Subtropical Fruits. Macmillan Co. pp.272-283. (Facsimile reprint 1974, Hafner Press, New York).
- Schroeder, C.A. 1946. Priority of the Species *Psidium cattleianum* Sabine. Jour. Arnold Arboretum. 27:314-315.
- Small, J.K. 1933. Manual of the Southeastern Flora. Univ. of No. Carolina Press, Chapel Hill. p. 937.
- The Staff of the Liberty Hyde Bailey Hortorium. 1976. Hortus Third. Macmillan Publishing Co., New York. p. 923.
- Stressau, EB. 1986. Florida, MyEden. Florida Classics Library, Port Salerno, FL. p. 102.

- Wunderlin, R.P. 1998. Guide to the Vascular Plants of Florida. Univ. Press of Florida, Gainesville. p. 454.
- Wunderlin, R.P., B.E Hansen, and E.L. Bridges. 1995 (updated May 1996). Atlas of Florida Vascular Plants. Web site: <http://www.usf.edu/~isb/projects/hb-atlas.html>.

Wildland WEEDS

Charter Advertisers

SePRO
Griffin
Dow Agrosciences
Riverdale
Invasive Plant Control
Brewer International
Timberland Enterprises, Inc.
American Cyanamid
Helena

New Name! Growing Commitment!

PROSOURCE ONE

AgroDistribution LLC. dba Prosource One

ProSource One formerly Terra Professional Products has a new name with a growing commitment to our customers.

ProSource One is the exclusive source for all of your aquatic vegetation management needs. We offer the right products, reliable advice and dependable services to help make your aquatic program successful. Talk to your ProSource One aquatic vegetation management specialist.

Aquatic Specialists
Western Florida
Eastern Florida

Polly Ellinor
Paul Mason

1-888-813-0562
1-800-207-1408